

# 2019 Black Shank Fungicide Trial

## Kent Boyd Farm – Hopkinsville KY

### Effect of Fungicides and Fumigation on Black Shank

- KT D6LC
  - Race 0 = 3; Race 1 = 3
  - Transplanted June 12
  - 4916 plants/A
- Fumigation used in 2 trts
  - Chloropicrin banded under row
  - 3 weeks prior to transplanted


Treatment	Rate/A	Timing
Orondis/Ridomil No Fumigation	4.8 oz Oron/8 oz Rid	TPW
Orondis/Ridomil Presidio No Fumigation	4.8 oz Oron/8 oz Rid 4 oz/A	TPW 1 <sup>st</sup> Cult
Orondis/Ridomil Presidio Ridomil No Fumigation	4.8 oz Oron/8 oz Rid 4 oz/A 16 oz/a	TPW 1 <sup>st</sup> Cult Layby
Orondis/Ridomil Presidio Ridomil Fumigated	4.8 oz Oron/8 oz Rid 4 oz/A 16 oz/a	TPW 1 <sup>st</sup> Cult Layby
No Fungicides Fumigated	-	-

# 2019 Black Shank Fungicide Trial

## Kent Boyd Farm – Hopkinsville KY

### Effect of Fungicides and Fumigation on Black Shank

LSD<sub>0.10</sub> = 620  
 ■ Total Yield



Final % Stand: 88.2 a  
 (LSD<sub>0.10</sub> = 20.5)

87.1 a

89.4 a

95.7 a

66.0 b